

Queensbury School
Wood End Road
Erdington
Birmingham
B24 8BL
Telephone: 0121 373 5731
Email: enquiry@queensbury.bham.sch.uk
Website: <https://www.queensburysch.com>



6th December 2019

Dear Parent/Carer

This letter is to advise you that diarrhoea and vomiting illness, which is present in the community at this time of the year, is also circulating at school.

We are following advice from Public Health England (PHE) West Midlands on managing the outbreak, including increasing cleaning throughout the school.

We would be very grateful for the support of parents in helping us reduce the spread of the infection. If your child has symptoms of diarrhoea and or vomiting, please keep them at home until 48 hours after they are symptom free. During this time, they should not mix with other children outside the home or visit local venues.

Norovirus is commonly known as Winter Vomiting Disease and often occurs in the community and in schools and nurseries at this time of the year. If your child is ill, you may find the following information useful:

ADVICE FOR PARENTS ABOUT NOROVIRUS

What are the symptoms?

- diarrhoea and vomiting usually lasting 1 -2 days

What should you do if your child is unwell?

- Make sure they get lots of rest.
- Ensure they drink plenty of fluids, taking sips rather than gulps to avoid vomiting.
- Give infant Paracetamol or Ibuprofen, according to product instructions, to help keep their temperature down.
- If your child is unusually sleepy, won't take fluids or has other symptoms, such as blood in their diarrhoea, an unusual rash, headache, neck stiffness or difficulty breathing, ring NHS Direct (0845 4647), contact your GP or take your child to hospital.
- Please ring your child's nursery/ school and let them know that your child is ill and what symptoms they have.



Interim Head Teacher: Mr Chris Wilson BSc/MA



Queensbury School

Wood End Road

Erdington

Birmingham

B24 8BL

Telephone: 0121 373 5731

Email: enquiry@queensbury.bham.sch.uk

Website: <https://www.queensburysch.com>



- Keep them at home until they are well enough to return to school. However, if your child has been suffering from diarrhoea and vomiting keep them at home until they have been free of symptoms for 48 hours.

How can you prevent the spread of these infections?

Hand washing is one of the most important ways to prevent the spread of both of these infections. This applies to the child who is ill and the person looking after them.

Hands should always be washed, using liquid soap if possible:

- Before and after caring for your child
- After using the toilet
- Before eating or handling food
- After cleaning up a mess (vomit, faeces or urine)

Other ways of preventing the spread of infection are to:

- Keep a separate towel for family members who have symptoms
- Dispose promptly of used tissues and other items that your child may have coughed or sneezed on
- Wash soiled clothing, bed linen and towels in a washing machine. Use the hottest wash for that fabric
- Clean baths and washbasins thoroughly and disinfect after use
- If cleaning up diarrhoea or vomit, wash the surface with hot soapy water and disinfect before allowing to dry. If using bleach remember that it can remove the colour from fabrics and can burn the skin

Please keep your child away from other children and people that are particularly vulnerable, eg the elderly and those with chronic illnesses.

Yours sincerely

Mr C D Wilson

Interim Head Teacher



Interim Head Teacher: Mr Chris Wilson BSc/MA





Winter Infections in Children **An information leaflet for parents**

This leaflet gives advice on illnesses that are currently affecting nursery and primary school children across the West Midlands

What are the symptoms?

There are two groups of symptoms

- Diarrhoea & vomiting usually lasting 1 - 2 days
- Fever, tiredness and nausea (feeling sick) sore throat, sometimes with a cough lasting 3 - 4 days

What should you do if your child is unwell?

- Keep your child at home until they are well

If your child has diarrhoea and/or vomiting they should not attend school/nursery until they have been free of symptoms for 48 hours

- Ensure they drink plenty of fluids, taking sips rather than gulps to avoid vomiting
- If they have a fever, give infant paracetamol and/or ibuprofen according to manufacturer's instructions
- If your child is unusually sleepy, is unable to take fluids or has other symptoms such as an unusual rash, headache or neck stiffness seek medical advice immediately.

Infection Control Advice

Hand washing is one of the most important ways to prevent the spread of both of these infections. This applies to the child who is ill and the person caring for them. Hands should be washed using soap and rinsed under running water:

- Before and after caring for child
- After using the toilet
- Before eating, preparing or handling food
- After cleaning up spills (vomit, diarrhoea or urine)
- Keep a separate towel for each family member who has symptoms. These should be changed regularly
- Dry your hands thoroughly

Other control measures include

- Dispose of used tissues immediately after use
- Wash soiled clothing , bed linen and towels in a washing machine, using the hottest wash for the fabric
- Ensure toilet areas, baths, and washbasins are kept clean using a solution of hot soapy water
- After initial cleaning, areas may also be disinfected using a 1 in 10 bleach solution. *(If using bleach remember that it can remove the colour from fabrics and can burn the skin)*

Further information can be obtained from:

West Midlands Public Health England team

2nd Floor, Kidderminster Library, Market Street, Kidderminster, DY10 1AB